

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

August 11, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/496,738
FILING DATE: August 21, 2003

Certified by

Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office



BEST AVAILABLE COPY

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. 60/496138

INVENTOR(S)			
Given Name (first and middle (if any))	Family Name or Surname	Residence (City and either State or Foreign Country)	
Michael Sandra Jorn	Didriksen Hogg Willigers Arnt	Vipperod, Denmark Bagsværd, Denmark Solrod Strand, Denmark	
<i>Additional inventors are being named on the _____ separately numbered sheets attached hereto</i>			
TITLE OF THE INVENTION (500 characters max)			
THE COMBINATION OF A SEROTONIN REUPTAKE INHIBITOR AND A GLYCINE TRANSPORTER TYPE 1 INHIBITOR FOR THE TREATMENT OF DEPRESSION			
Direct all correspondence to: CORRESPONDENCE ADDRESS			
<input checked="" type="checkbox"/> Customer Number: 07278 <small>Customer Number</small>			
OR			
<input checked="" type="checkbox"/> Firm or Individual Name S. Peter Ludwig DARBY & DARBY P.C.			
Address P.O. Box 5257			
City New York		State NY	
Country US		Zip 10150-5257	
		Telephone (212) 527-7700	
		Fax (212) 753-6237	
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification Number of Pages 34		<input type="checkbox"/> CD(s), Number 	
<input type="checkbox"/> Drawing(s) Number of Sheets 		<input type="checkbox"/> Other 	
<input checked="" type="checkbox"/> Application Data Sheet. See 37 CFR 1.76 (specify): 			
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT			
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.			
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fees.			
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number:		04-0100	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.		FILING FEE AMOUNT (\$) 160.00	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.			
<input checked="" type="checkbox"/> No <input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: 			

[Page 1 of 1]

Respectfully submitted,

SIGNATURE

J.P. Lesser

TYPED OR

PRINTED NAME

Jay P. Lesser

TELEPHONE

(212) 527-7770

Date August 21, 2003

REGISTRATION NO.

41,151

(if appropriate)

Docket Number:

5432/0N212US0

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

εJ306629631-us

Express Mail Label No.

Dated: 8/21/03

B.W. Lee



Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PTO/SB/17 (10-02)

Approved for use through 10/31/2002, OMB 0951-0032

FEE TRANSMITTAL for FY 2003

Patent fees are subject to annual revision.

☐ Applicant claims small entity status. See 37 CFR 1.27

TOTAL AMOUNT OF PAYMENT (\$)

160.00

Complete if Known

Application Number Not Yet Assigned
Filing Date
First Named Inventor Michael Didriksen
Examiner Name Not Yet Assigned
Group Art Unit N/A
Attorney Docket No. 5432/0N212US0

METHOD OF PAYMENT (check all that apply)

☒ Check ☐ Credit Card ☐ Money Order ☐ Other ☐ None

☐ Deposit Account

Deposit Account Number

Deposit Account Name

The Commissioner is hereby authorized to: (check all that apply)

☐ Charge fee(s) indicated below ☒ Credit any overpayments

☐ Charge any additional fee(s) during the pendency of this application

☐ Charge fee(s) indicated below, except for the filing fee

to the above-identified deposit account.

FEE CALCULATION

1. BASIC FILING FEE

Large Entity	Small Entity	Fee Code	Fee (\$)	Fee Description	Fee Paid
		1001	740	Utility filing fee	
		1002	330	Design filing fee	
		1003	510	Plant filing fee	
		1004	740	Reissue filing fee	
		1005	160	Provisional filing fee	160.00
SUBTOTAL (1) (\$)					160.00

2. EXTRA CLAIM FEES FOR UTILITY AND REISSUE

Total Claims	Extra Claims	Fee from below	Fee Paid
Independent Claims	x	=	
Multiple Dependent	x	=	

Large Entity	Small Entity	Fee Code	Fee (\$)	Fee Description	Fee Paid
		1202	18	Claims in excess of 20	
		1201	84	Independent claims in excess of 3	
		1203	280	Multiple dependent claim, if not paid	
		1204	84	Reissue independent claims over original patent	
		1205	18	Reissue claims in excess of 20 and over original patent	
SUBTOTAL (2) (\$)					0.00

**or number previously paid, if greater; For Reissues, see above

FEE CALCULATION (continued)

3. ADDITIONAL FEES

Large Entity	Small Entity	Fee Code	Fee (\$)	Fee Description	Fee Paid
		1051	130	Surcharge - late filing fee or oath	
		1052	50	Surcharge - late provisional filing fee or cover sheet	
		1053	130	Non-English specification	
		1812	2,520	For filing a request for ex parte reexamination	
		1804	920*	Requesting publication of SIR prior to Examiner action	
		1805	1,840*	Requesting publication of SIR after Examiner action	
		1251	110	Extension for reply within first month	
		1252	400	Extension for reply within second month	
		1253	920	Extension for reply within third month	
		1254	1,440	Extension for reply within fourth month	
		1255	1,960	Extension for reply within fifth month	
		1401	320	Notice of Appeal	
		1402	320	Filing a brief in support of an appeal	
		1403	280	Request for oral hearing	
		1451	1,510	Petition to institute a public use proceeding	
		1452	110	Petition to revive - unavoidable	
		1453	1,280	Petition to revive - unintentional	
		1501	1,280	Utility issue fee (or reissue)	
		1502	460	Design issue fee	
		1503	620	Plant issue fee	
		1460	130	Petitions to the Commissioner	
		1807	50	Processing fee under 37 CFR 1.17(c)	
		1806	180	Submission of Information Disclosure Sheet	
		8021	40	Recording each patent assignment per property (times number of properties)	
		1809	740	Filing a submission after final rejection (37 CFR 1.129(a))	
		1810	740	For each additional invention to be examined (37 CFR 1.129(b))	
		1801	740	Request for Continued Examination (RCE)	
		1802	900	Request for expedited examination of a design application	
Other fee (specify)					

*Reduced by Basic Filing Fee Paid

SUBTOTAL (3) (\$)

0.00

SUBMITTED BY

Name (Print/Type)

Signature

Jay P. Lessler

Registration No. (Attorney/Agent)

41,151

Complete (if applicable)

Telephone (212) 527-7765

Date

August 21, 2003

Application No. (if known):

Attorney Docket No.: 5432/0N212US0

Certificate of Express Mailing Under 37 CFR 1.10

I hereby certify that this correspondence is being deposited with the United States Postal Service as
Express Mail, Airbill No. _____ in an envelope addressed to:

MS Provisional Patent Application **EW 306629631-US**
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

on August 21, 2003
Date

B.W. Lee

Signature

B.W. LEE

Typed or printed name of person signing Certificate

Note: Each paper must have its own certificate of mailing, or this certificate must identify each submitted paper.

Provisional Application Transmittal (1 page)
Specification and Claims (34 pages)
Application Data Sheet (3 pages)
Fee Transmittal (1 page)
Check No. 2379 in the amount of \$160.00

Application Data Sheet

Application Information

Application Type::	Provisional
Subject Matter::	Utility
Suggested Group Art Unit::	N/A
CD-ROM or CD-R?::	None
Sequence submission?::	None
Computer Readable Form (CRF)?::	No
Title::	THE COMBINATION OF A SEROTONIN REUPTAKE INHIBITOR AND A GLYCINE TRANSPORTER TYPE 1 INHIBITOR FOR THE TREATMENT OF DEPRESSION
Attorney Docket Number::	5432/0N212US0
Request for Early Publication?::	No
Request for Non-Publication?::	No
Small Entity?::	No
Petition included?::	No
Secrecy Order in Parent Appl.?::	No

Applicant Information

Applicant Authority Type::	Inventor
Primary Citizenship Country::	Denmark
Status::	Full Capacity
Given Name::	Michael
Family Name::	Didriksen
City of Residence::	Vipperod
Country of Residence::	Denmark
Street of mailing address::	Kilevangsparken 66
City of mailing address::	Vipperod
Country of mailing address::	Denmark
Postal or Zip Code of mailing address::	4390

Applicant Authority Type:: Inventor
Primary Citizenship Country:: British
Status:: Full Capacity
Given Name:: Sandra
Family Name:: Hogg Willigers
City of Residence:: Bagsværd
Country of Residence:: Denmark
Street of mailing address:: Helmsvej 38
City of mailing address:: Bagsværd
Country of mailing address:: Denmark
Postal or Zip Code of mailing address:: 2880

Applicant Authority Type:: Inventor
Primary Citizenship Country:: Denmark
Status:: Full Capacity
Given Name:: Jorn
Family Name:: Arnt
City of Residence:: Solrod Strand
Country of Residence:: Denmark
Street of mailing address:: Svend Gonges Vej 11 A
City of mailing address:: Solrod Strand
Country of mailing address:: Denmark
Postal or Zip Code of mailing address:: 2680

Correspondence Information

Correspondence Customer Number:: 07278

Representative Information

Representative Customer Number:: 07278

Assignee Information

Assignee name::	H. Lundbeck A/S
Street of mailing address::	Ottiliavej 9
City of mailing address::	Copenhagen-Valby
Country of mailing address::	Denmark
Postal or Zip Code of mailing address::	DK-2500

The combination of a serotonin reuptake inhibitor and a glycine transporter type 1 inhibitor for the treatment of depression

The present invention relates to the combination of a serotonin reuptake inhibitor (SRI) and a glycine transporter type 1 (GlyT-1) inhibitor. Accordingly, the present invention relates to the use of certain compounds, and to compositions of compounds having serotonin reuptake inhibiting activity and GlyT-1 inhibitor activity for the treatment of depression and other affective disorders.

Background

Selective serotonin reuptake inhibitors (hereinafter referred to as SSRIs) have become first choice therapeutics in the treatment of depression, certain forms of anxiety and social phobias, because they are effective, well tolerated and have a favourable safety profile compared to the classic tricyclic antidepressants.

However, clinical studies on depression and anxiety disorders indicate that non-response to SSRIs is substantial, up to 30%. Another, often neglected, factor in antidepressant treatment is compliance, which has a rather profound effect on the patient's motivation to continue pharmacotherapy.

First of all, there is the delay in therapeutic effect of SSRIs. Sometimes symptoms even worsen during the first weeks of treatment. Secondly, sexual dysfunction is a side effect common to all SSRIs. Without addressing these problems, real progress in the pharmacotherapy of depression and anxiety disorders is not likely to happen.

In order to cope with non-response, psychiatrists sometimes make use of augmentation strategies. Augmentation of antidepressant therapy may be accomplished through the co-administration of mood stabilizers such as lithium carbonate or triiodothyronin or by the use of electroshock.

In 1993, an augmentation strategy with pindolol was described by Artigas et al. in *Trends Pharmacol. Sci.* 1993, 14, p 262-263. Artigas' idea is based on intracerebral

EXPRESS MAIL CERTIFICATE

Date 8/21/03 Label No. EV 306629631-45
I hereby certify that, on the date indicated above, this paper or
fee was deposited with the U.S. Postal Service & that it was
addressed for delivery to the Commissioner for Patents,
P.O. Box 1450, Alexandria, VA 22313-1450 by "Express Mail
Post Office to Addressee" service.

B.W. LEE

B.W. Lee

microdialysis experiments in animals. In fact, later neurochemical studies built on the desensitization hypothesis by Blier and co-workers stated that the delay in therapeutic effect of antidepressants is related to a gradual desensitization of 5-HT autoreceptors (Blier et al. *J. Clin. Psychopharmacol.* 1987, 7 suppl. 6, 24S-35S). A key point in their hypothesis is that the effects of SSRIs on the release-controlling somatodendritic autoreceptors (5-HT_{1A}) limit the release of 5-HT in terminal areas and thus the effect of 5-HT uptake inhibition in those regions. This is supported by microdialysis experiments in rats, showing that the increase in extracellular 5-HT elicited by a single dose of an SSRI is augmented by co-administration of a 5-HT_{1A} autoreceptor antagonist (Invernizzi et al. *Brain Res.* 1992, 584, p 322-324 and Hjorth, S., *J. Neurochem.* 1993, 60, p 776-779).

The effect of combined administration of a compound that inhibits serotonin reuptake and a 5-HT_{1A} receptor antagonist has been evaluated in several studies (Innis, R.B. et al. *Eur. J. Pharmacol.* 1987, 143, p. 1095-204 and Gartside, S.E., *Br. J. Pharmacol.* 1995, 115, p 1064-1070, Blier, P. et al. *Trends in Pharmacol. Science* 1994, 15, 220). In these studies it was found that 5-HT_{1A} receptor antagonists would abolish the initial brake on 5-HT neurotransmission induced by the serotonin reuptake inhibitors and thus produce an immediate boost of 5-HT transmission and a rapid onset of therapeutic action.

Several patent applications have been filed which cover the use of a combination of a 5-HT_{1A} antagonist and a serotonin reuptake inhibitor for the treatment of depression (see e.g. EP-A2-687 472 and EP-A2-714 663).

Another approach to increase terminal 5-HT would be through blockade of the 5-HT_{1B} autoreceptor. Microdialysis experiments in rats have indeed shown that increase of hippocampal 5-HT by citalopram is potentiated by GMC 2-29, an experimental 5-HT_{1B} receptor antagonist.

Several patent applications covering the combination of an SSRI and a 5-HT_{1B} antagonist or partial agonist have also been filed (WO 97/28141, WO 96/03400, EP-A-701819 and WO 99/13877).

Glutamate is the most important excitatory neurotransmitter in the brain mediating its effect via ionotropic and metabotropic receptors. Ionotropic NMDA receptors are involved in the glutamatergic excitation of GABAergic, serotonergic, dopaminergic, and adrenergic neurons.

The NMDA receptor is positive modulated by glycine. Functional NMDA receptor complexes are formed by combinations of NR1 and NR2 subunits, which contain the glycine and glutamate recognition sites, respectively (Danysz W & Parsons C.G., Pharmacological reviews, vol 50: pp597-664 (1998)).

GlyT-1 transporters located in the adjacent glia cells regulate the endogenous level of glycine in the vicinity of the NMDA receptor complex. Consequently, inhibiting the GlyT-1 transporter results in increased level of glycine and NMDA receptor activation (Danysz W & Parsons C.G., Pharmacological reviews, vol 50: pp597-664 (1998)).

In preclinical models of depression (Chronic severe stress and Chronic mild stress) the involvement of the NMDA receptor complex has been shown (Novak G. et al., Polish Journal of Pharmacology, vol 58: pp365-369 (1998)). Further, Glycine site partial agonists show antidepressant like effect in the Chronic mild stress model (Papp M. & Moryl E., European Journal of Pharmacology, vol 316: pp145-151 (1996)).

Description of the invention

It has now surprisingly been found that a GlyT-1 inhibitor will augment the effect of an SRI, in particular an SSRI on extracellular 5-HT levels.

It is therefore suggested that the combination of an SSRI and a GlyT-1 inhibitor, provide 5-HT reuptake inhibitory and GlyT-1 inhibitor properties, and would have a better efficacy and faster onset than an SSRI alone.

The present invention thus provides:

The use of a GlyT-1 inhibitor for the preparation of a pharmaceutical composition to be used in combination with a serotonin reuptake inhibitor (SRI).

The present invention relates to the use of a compound, which is a serotonin reuptake inhibitor, and another compound, which is a GlyT-1 inhibitor for the preparation of a pharmaceutical composition for the treatment of depression, anxiety disorders and other affective disorders, such as generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder and social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse or any other disorder responsive to serotonin reuptake inhibitors.

The present invention also relates to the use of a GlyT-1 inhibitor for the preparation of a pharmaceutical composition useful for augmenting and providing faster onset of the therapeutic effect of a serotonin reuptake inhibitor. Moreover, the present invention also relates to the use of a GlyT-1 inhibitor for the preparation of a pharmaceutical composition useful for augmenting or providing faster onset of the therapeutic effect of a serotonin reuptake inhibitor.

Moreover the invention relates to the use of a combination of a compound, which is a serotonin reuptake inhibitor, and a compound, which is a GlyT-1 inhibitor, for the preparation of a pharmaceutical composition or kit-of-parts (kit) useful for the treatment of depression, anxiety disorders and other affective disorders, such as generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder and social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse or any other disorder responsive to serotonin reuptake inhibitors.

Furthermore the invention relates to the use of a compound, which is a serotonin reuptake inhibitor, and a compound, which is a GlyT-1 inhibitor, for the preparation

of a kit for use in the treatment of depression, anxiety disorders and other affective disorders, such as generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder and social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse or any other disorder responsive to serotonin reuptake inhibitors.

In a further aspect the invention relates to a pharmaceutical composition comprising a combination of a compound, which is a serotonin reuptake inhibitor, and another compound, which is a GlyT-1 inhibitor, and optionally pharmaceutically acceptable carriers or diluents.

In a further aspect the invention relates to a kit comprising a combination of a compound, which is a serotonin reuptake inhibitor, and another compound, which is a GlyT-1 inhibitor, and optionally pharmaceutically acceptable carriers or diluents.

In yet another aspect the invention relates to a method for the treatment of depression, anxiety disorders and other affective disorders, such as generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder and social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse or any other disorder responsive to serotonin reuptake inhibitors comprising administering to a person in need thereof a therapeutically effective amount of a combination of a compound, which is a serotonin reuptake inhibitor and a compound, which is a GlyT-1 inhibitor.

Each of the medical indications: depression, anxiety disorders and other affective disorders, including generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder or social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia,

premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse and any other disorder responsive to a SRI is intended to be an individual embodiment. Accordingly, whenever mentioned in the present description, each of the indications specified above may be claimed individually.

Whenever the indications depression, anxiety disorders and other affective disorders, including generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder or social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse and any other disorder responsive to a SRI are mentioned in relation to use of a GlyT-1 inhibitor and a SRI, a pharmaceutical composition, a kit, a method of treatment and a method for the identification of compounds useful for treatment each indication is intended to be an individual embodiment. Accordingly, each of the indications specified above may individually be claimed together with said use of a GlyT-1 inhibitor and an SRI, pharmaceutical composition, kit, method of treatment and method for the identification of compounds useful for treatment.

In a particular embodiment, the SRI is a selective serotonin reuptake inhibitor (SSRI).

In another particular embodiment, a GlyT-1 inhibitor, which is selective for the glycine transporter type 1 is used according to the invention.

The pharmaceutical composition or kit according to the invention may be administered by simultaneous administration. The term "simultaneous administration" as used herein means, that the GlyT-1 inhibitor and the SRI are administered with a time separation of no more than 15 minutes, such as at most 10 minutes, such as at most 5 minutes or such as at most 2 minutes. The GlyT-1 inhibitor and the SRI may be contained in the "same unit dosage form" or in "discrete dosage forms". As used herein, the term "same unit dosage form" means a dosage form comprising both the SRI and the GlyT-1 inhibitor. As used herein, the term "discrete dosage form" means

that the GlyT-1 inhibitor is comprised in one dosage form and that the SRI is comprised in another dosage form.

Simultaneous administration of the GlyT-1 inhibitor and the SRI is optionally combined with administration of supplementary doses of GlyT-1 inhibitor. The supplementary doses of GlyT-1 inhibitor may be given for instance 1, 2, 3 or 4 times a day whereas the SRI and the GlyT-1 inhibitor which are administered by "simultaneous administration" may be given one or more times a day, e.g. once daily or e.g. twice daily. Accordingly:

a) the GlyT-1 inhibitor and the SRI may be administered by simultaneous administration once daily and supplementary doses of GlyT-1 inhibitor may be administered 1, 2, 3 or 4 times a day, such as 1, 2 or 3 times a day, such as once or twice daily, such as twice daily or such as once daily,

or

b) the GlyT-1 inhibitor and the SRI may be administered by simultaneous administration twice daily and supplementary doses of GlyT-1 inhibitor may be administered 1, 2, 3 or 4 times a day, such as 1, 2 or 3 times a day, such as once or twice daily, such as twice daily or such as once daily.

Alternatively, the pharmaceutical composition or kit according to the invention is administered by sequential administration. The term "sequential administration" as used herein means that one (1) or more daily doses of the GlyT-1 inhibitor and 1 or more daily doses of SRI are administered with a time separation between two administered doses of more than 15 minutes and less than 4 hours, such as more than 2 hours and less than 4 hours, such as more than 15 minutes and less than 2 hours, such as more than 1 hour and less than 2 hours, such as more than 30 minutes and less than 1 hour, such as more than 15 minutes and less than 30 minutes. Either the SRI or the GlyT-1 inhibitor may be administered first. The GlyT-1 inhibitor and the SRI are contained in discrete dosage forms, optionally contained in the same container or package. Typically, 1, 2, 3, 4 or 5 daily doses of GlyT-1 inhibitor and 1 or 2 daily doses of SRI may be administered. Accordingly:

a) the GlyT-1 inhibitor and the SRI may be administered once daily and the GlyT-1 inhibitor may be administered 1, 2, 3, 4 or 5 times a day, such as 1, 2, 3 or 4 times a

day, such as 1, 2 or 3 times a day, such as once or twice daily, such as twice daily or such as once daily,

or

b) the GlyT-1 inhibitor and the SRI may be administered twice daily and the GlyT-1 inhibitor may be administered 1, 2, 3, 4 or 5 times a day, such as 1, 2, 3 or 4 times a day, such as 1, 2 or 3 times a day, such as once or twice daily, such as twice daily or such as once daily.

Accordingly, the pharmaceutical composition or kit according to the invention may be adapted for simultaneous administration of the active ingredients, or it may be adapted for sequential administration of the active ingredients. When the pharmaceutical composition or kit is adapted for simultaneous administration, the active ingredients may be contained in the same unit dosage form. When the pharmaceutical composition or kit is adapted for sequential administration, the active ingredients are contained in discrete dosage forms, optionally contained in the same container or package. As used herein, an "active ingredient" means a SRI or a GlyT-1 inhibitor.

A kit comprises a preparation of the GlyT-1 inhibitor in a first-unit dosage form, and the SRI in a second-unit dosage form, and container means for containing said first and second dosage forms.

In a further embodiment the GlyT-1 inhibitor is selected from any one of the compounds disclosed in WO0208216, such as any one of

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl} glycine ethyl ester,

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine ethyl ester,

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl} glycine,

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[1-(3-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[5-trifluoromethyl-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylalanine,
 N-{3-[5-cyano-1-(3-methyl-4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-cyano-1-(4-cyanophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylalanine,
 N-{3-[5-cyano-1-(4-methoxyphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine, N-{2-[5-cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]ethyl}-N-methylglycine,
 N-{3-[5-Chloro-1-(4-chloro-phenyl)-indan-1-yl]-propyl}-N-methylglycine,
 N-{3-[5-Chloro-1-(4-chloro-phenyl)-indan-1-yl]-propyl}-N-methylalanine,
 N-{3-[3-cyclo-1-(4-methylphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-benzo[c]thiophen-1-yl)-propyl}-N-methylglycine,
 N-{3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-benzo[c]thiophen-1-yl)-propyl}-N-methylalanine,
 N-{3-[1-(4-Fluoro-phenyl)-3,3-dimethyl-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methylglycine,
 N-{3-[5-Bromo-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{2-[1-(4-Chloro-phenyl)-3,3-dimethyl-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methylglycine,
 N-[3-(3-methyl-1-phenyl-1*H*-inden-1-yl)-propyl]-N-methylglycine,
 N-[3-(5-Chloro-1-thiophen-2-yl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylglycine,
 N-[3-(5-Chloro-1-thiophen-2-yl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methyl(1-ethyl)-glycine,
 N-[3-(3-methyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylalanine,
 N-[3-(3-methyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methyl(1-ethyl)-glycine,

N-[3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methylalanine,
 N-[3-(3,3-Dimethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methylalanine,
 N-[3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methyl-(1-ethyl)glycine,
 N-[3-(3,3-Dimethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methyl-(1-ethyl)glycine,
 N-[3-(3,3-Diethyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylalanine,
 N-[3-(3,3-Diethyl-1-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylalanine,
 N-[3-(3,3-Diethyl-1-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylglycine,
 N-[3-(1-phenyl-1,3-dihydro-benzo[c]thiophen-1-yl)-propyl]-N-methylalanine,
 N-{3-[1-(4-Chloro-phenyl)-3,3-dimethyl-indan-1-yl]-propyl}-N-methylglycine,
 N-{3-[1-(4-Chloro-phenyl)-3,3-diethyl-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-alanine,
 N-[2-(3-methyl-1-phenyl-indan-1-yl)-ethyl]-amino)-N-methyl alanine,
 N-[3-(1-phenyl-(1*H*)-inden-1-yl)-propyl]-N-methyl-alanine,
 N-{3-[1-(4-Fluoro-phenyl)-5-(4-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[5-Chloro-1-(4-chloro-phenyl)-indan-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[5-Chloro-1-(4-chloro-phenyl)-indan-1-yl]-propyl}-N-methyl-alanine,
 N-{3-[1-(4-chloro-phenyl)-5-(4-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(4-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(4-methoxy-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2-thiophenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,

N-{3-[1-(4-Chloro-phenyl)-5-(4-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(4-methoxy-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-chloro-phenyl)-5-(4-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{2-[1-(4-Chloro-phenyl)-5-(5-chloro-thiophen-2-yl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(3-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2,5-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-chloro-phenyl)-5-(3-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-chloro-phenyl)-5-(3-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine ,
 N-{3-[1-(4-Chloro-phenyl)-5-(3,4-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(3-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2,5-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(3,4-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,

N-{3-[1-(4-chloro-phenyl)-5-(2-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine ,
or a pharmaceutically acceptable addition salt thereof.

In a further embodiment the GlyT-1 inhibitor is selected from any one of the compounds disclosed in WO03/053942, such as any one of

(+/-)-{4-[2-(4-Methoxy-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-Chloro-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-Fluoro-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-*iso*-Propyl-phenylsulfanyl)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
(+/-)-2-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethylpiperazin-1-yl}-propionic acid,
{4-[5-Chloro-2-(4-methoxy-phenylsulfanyl)-phenyl]-2(R)-methyl-piperazin-1-yl}-acetic acid,
{4-[2-(4-Methoxy-phenylsulfanyl)-phenyl]-2(R),5(S)-dimethyl-piperazin-1-yl}-acetic acid,
{4-[5-Chloro-2-(4-methoxy-phenylsulfanyl)-phenyl]-2,2-dimethyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[5-Chloro-2-(4-trifluoromethyl-phenylsulfanyl)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
{4-[5-Chloro-2-(3-methoxy-phenylsulfanyl)-phenyl]-2(R)-methyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-Phenyl-phenyloxy)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
(+/-)-{4-[2-(4-Methyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,

(+/-)-{4-[2-(4-*iso*-Propyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(2,4-Dimethyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-2-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-3-methylpiperazin-1-yl}-propionic acid,
 {4-[2-(4-Isopropyl-phenylsulfanyl)-phenyl]-piperazin-1-yl}-acetic acid,
 (+/-)-2-{4-[2-(4-Methoxy-phenylsulfanyl)-phenyl]-3-methyl-piperazin-1-yl}-propionic acid,
 or a pharmaceutically acceptable acid addition salt thereof.

Typical GlyT-1 inhibitors show inhibition below 20000 nM as IC₅₀ in the "[³H]-Glycine uptake" test described herein.

The invention also covers GlyT-1 inhibitors identified according to this method, but is not limited to these assay methods.

According to the invention, it has been found that co-administration of GlyT-1 inhibitors and a serotonin reuptake inhibitor produces a significant increase in the level of serotonin in terminal areas, as measured in microdialysis experiments, compared to the administration of the serotonin reuptake inhibitor alone.

According to the invention, animal studies have shown that GlyT-1 inhibitors may provide fast onset of therapeutic effect of serotonin reuptake inhibitors and potentiate the anxiolytic potential of serotonin reuptake inhibitors.


The use of a combination of a GlyT-1 inhibitor and a serotonin reuptake inhibitor may greatly reduce the amount of serotonin reuptake inhibitor necessary to treat depression and other affective disorders and may thus reduce the side effects caused by the serotonin reuptake inhibitor. In particular, the combination of a reduced amount of SRI and a GlyT-1 inhibitor may reduce the risk of SSRI-induced sexual dysfunction and sleep disturbances.

Co-administration of a GlyT-1 inhibitor and a serotonin reuptake inhibitor may also be useful for the treatment of refractory depression, i.e. depression, which cannot be treated appropriately by administration of a serotonin reuptake inhibitor alone. Typically, GlyT-1 inhibitors may be used as add-on therapy for the augmentation of the response to SRI in patients where at least 40-60% reduction in symptoms has not been achieved during the first 6 weeks of treatment with an SRI.

Many antidepressants with serotonin reuptake inhibiting effect have been described in the literature. Any pharmacologically active compound, which primarily or partly exert its therapeutic effect via inhibition of serotonin reuptake in the CNS, may benefit from augmentation with a GlyT-1 inhibitor.


The following list contains a number of serotonin reuptake inhibitors, which may benefit from augmentation with a GlyT-1 inhibitor: citalopram, escitalopram, fluoxetine, R-fluoxetine, sertraline, paroxetine, fluvoxamine, venlafaxine, desmethylvenlafaxine, duloxetine, dapoxetine, vilazodone, nefazodone, imipramine, imipramine N-oxide, desipramine, pirandamine, dazepinil, nefopam, befuraline, fezolamine, femoxetine, clomipramine, cianoimipramine, litoxetine, cericlamine, seproxetine, WY 27587, WY 27866, imeldine, ifoxetine, indeloxazine, tiflucarbine, viqualine, milnacipran, bazinaprine, YM 922, S 33005, F 98214-TA, FI 4503, A 80426, EMD 86006, NS 2389, S33005, OPC 14523, alaproclate, cyanodothepine, trimipramine, quinupramine, dothiepin, amoxapine, nitroxazepine, McN 5652, McN 5707, VN 2222, L 792339, roxindole, YM 35992, OI 77, Org 6582, Org 6997, Org 6906, amitriptyline, amitriptyline N-oxide, nortriptyline, CL 255.663, pirlindole, indatraline, LY 280253, LY 285974, LY 113.821, LY 214.281, CGP 6085 A, RU 25.591, napamezole, diclofensine, trazodone, BMY 42.569, NS 2389, sercloremin, nitroquipazine, ademethionine, sibutramine, desmethylsubitramine, didesmethylsubitramine, clovoxamine vilazodone. The compounds mentioned above may be used in the form of the base or a pharmaceutically acceptable acid addition salt thereof. Each of the serotonin reuptake inhibitors specified above is intended to be an individual embodiment. Accordingly, each of them and the use thereof may be claimed individually.

Compounds such as citalopram, escitalopram, fluoxetine, R-fluoxetine, sertraline, paroxetine, fluvoxamine, venlafaxine, desmethylvenlafaxine, duloxetine, dapoxetine, vilazodone, nefazodone, imipramine, imipramine N-oxide, desipramine, pirandamine, dazepinil, nefopam, befuraline, fezolamine, femoxetine, clomipramine, cianoimipramine, litoxetine, cericlamine, seproxetine, imeldine, ifoxetine, indeloxazine, tiflucarbine, viqualine, milnacipran, bazinaprine, alaproclate, cyanodothepine, trimipramine, quinupramine, dothiepin, amoxapine, nitroxazepine, roxindole, amitriptyline, amitriptyline N-oxide, nortriptyline, pirlindole, indatraline, napamezole, diclofensine, trazodone, sercloremin, nitroquipazine, ademethionine, sibutramine, desmethylsibutramine, didesmethylsibutramine, clovoxamine vilazodone, N-[(1-[(6-Fluoro-2-naphthalenyl)methyl]-4-piperidinyl)amino]carbonyl]-3-pyridine carboxamide (WY 27587), [trans-6-(2-chlorophenyl)-1,2,3,5,6,10b-hexahydropyrrolo-(2,1-a)isoquinoline] (McN 5707), (dl-4-exo-amino-8-chloro-benzo-(b)-bicyclo[3.3.1]nona-2-6 alpha(10 alpha)-diene hydrochloride)(Org 6997), (dl)-(5 alpha,8 alpha,9 alpha)-5,8,9,10-Tetrahydro-5,9- methanobenzocycloocten-8-amine hydrochloride (Org 6906), -[2-[4-(6-fluoro-1H-indol-3-yl)-3,6-dihydro-1(2H)-pyridinyl]ethyl]-3-isopropyl-6-(methylsulphonyl)-3,4-dihydro-1H-2,1,3-benzothiadiazine-2,2-dioxide (LY393558), [4-(5,6-dimethyl-2-benzofuranyl)-piperidine] (CGP 6085), dimethyl-[5-(4-nitro-phenoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-7-yl]-amine (RU 25.591),




(A 80426),


17




(EMD 86006),





(S33005),



(OPC 14523),



(McN 5652),



are preferred. The compounds mentioned above may be used in the form of the base or a pharmaceutically acceptable acid addition salt thereof. Each of the serotonin reuptake inhibitors specified above is intended to be an individual embodiment. Accordingly, each of them and the use thereof may be claimed individually.

In a further embodiment the SRI is selected from citalopram, escitalopram, fluoxetine, sertraline, paroxetine, fluvoxamine, venlafaxine, dapoxetine, vilazodone, duloxetine, nefazodone, imipramine, femoxetine and clomipramine, preferably citalopram, or escitalopram.

Typical serotonin reuptake inhibitors show serotonin reuptake inhibition below 10000 nM (IC_{50}) in the "Inhibition of the uptake of [3H]Serotonin into whole rat brain synaptosomes" test described herein.

Other therapeutic compounds, which may benefit from augmentation with GlyT-1 inhibitors, include compounds, which cause an elevation in the extracellular level of

Accordingly, other compounds than SRIs which cause an elevation in the extracellular level of serotonin, may be used instead of SRIs in every aspect of the invention as described herein.

The above list of serotonin reuptake inhibitors and other compounds, which cause an increase in the extracellular level of serotonin, may not be construed as limiting.

The term selective serotonin reuptake inhibitor (SSRI) means an inhibitor of the monoamine transporters, which has stronger inhibitory effect at the serotonin transporter than the dopamine and the noradrenaline transporters. Particularly preferred SSRIs according to the invention are citalopram, escitalopram, fluoxetine, fluvoxamine, sertraline, duloxetine, vilazodone and paroxetine.

Each of the active ingredients according to the invention may be administered alone or together or in combination with pharmaceutically acceptable carriers or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19 Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the specific active ingredient or active ingredients chosen.

Pharmaceutical compositions for oral administration include solid dosage forms such as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, they may be prepared with coatings such as enteric coatings or they may be formulated so as to provide controlled release of one or more active ingredient such as sustained or prolonged release according to methods well known in the art.

Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

Pharmaceutical compositions for parenteral administration include sterile aqueous and nonaqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants etc.

The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally in the form of solutions for injection. For preparing such compositions, methods well known in the art may be used, and any pharmaceutically acceptable carriers, diluents, excipients or other additives normally used in the art may be used.

A typical oral dosage of each of the active ingredients is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the

condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is a base addition salt of a compound having the utility of a free acid. When an active ingredient contains a free acid such salts are prepared in a conventional manner by treating a solution or suspension of a free acid of the active ingredient with a chemical equivalent of a pharmaceutically acceptable base.

For parenteral administration, solutions of one or more active ingredient in sterile aqueous solution, aqueous propylene glycol, aqueous vitamin E or sesame or peanut oil may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

Solutions for injections may be prepared by dissolving one or more active ingredients and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to a desired volume, sterilising the solution and filling it in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents.

Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, agar, pectin, acacia, stearic acid and lower alkyl ethers of cellulose corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like.

Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredient or ingredients used.

Examples of liquid carriers are syrup, peanut oil, olive oil, phospho lipids, fatty acids, fatty acid amines, polyoxyethylene and water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

The pharmaceutical compositions formed by combining one or more active ingredients of the invention with the pharmaceutical acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

The active ingredients of the invention may be formulated in similar or dissimilar pharmaceutical compositions and unit forms thereof.

If a solid carrier is used for oral administration, the preparation may be tablette, placed in a hard gelatine capsule in powder or pellet form or it may be in the form of a troche or lozenge.

The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g.

If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

If desired, the pharmaceutical composition of the invention may comprise one or more active ingredients in combination with further pharmacologically active substances such as those described in the foregoing.

Materials and Methods

[³H]-Glycine uptake

The GlyT-1 inhibitors for use in combination with an SRI, such as an SSRI, are tested in the well-recognised and reliable test measuring glycine uptake:

Cells transfected with the human GlyT-1b were seeded in 96 well plates. Prior to the experiment the cells were washed twice in HBS (10 mM Hepes-tris (pH 7.4), 2,5 mM KCl, 1 mM CaCl₂, 2,5 mM MgSO₄) and pre-incubated with test compound for 6 minutes. Afterwards, 10 nM ³H-glycine was added to each well and the incubation was continued for 15 minutes. The cells were washed twice in HBS. Scintillation fluid was added and the Plates were counted on a Trilux (Wallac) scintillation counter.

Based on this test, compounds which are GlyT-1 inhibitors show inhibition below 20000 nM as IC₅₀ in the above-mentioned assay, preferably below 10000 nM.

Inhibition of the uptake of [³H]Serotonin into whole rat brain synaptosomes

The inhibition of the serotonin uptake of an SRI is tested in the well-recognised and reliable test measuring serotonin uptake:

The compounds were tested with respect to their 5-HT reuptake inhibiting effect by measuring their ability to inhibit the uptake of [³H]serotonin into whole rat brain synaptosomes *in vitro*. The assay was performed as described by Hyttel *Psychopharmacology* 1978, 60, 13.


Based on this test, compounds which are SRI exhibit serotonin reuptake inhibition below 10000 nM (IC₅₀) in the assay above.

Animals

Male albino rats of a Wistar-derived strain (285-320 g; Harlan, Zeist, The Netherlands) were used for the experiments. Upon surgery, rats were housed individually in plastic cages (35 x 35 x 40 cm), and had free access to food and water. Animals were kept on a 12 h light schedule (light on 7:00 a.m.). The experiments are concordant with the declarations of Helsinki and were approved by the animal care committee of the faculty of mathematics and natural science of the University of Groningen .

Drugs

The following drugs were used: Citalopram hydrobromide and NFPS having the structure:



(LU 2736N) (Lundbeck A/S, Copenhagen, Denmark). Drugs were dissolved in saline and administered s.c.

Surgery

Microdialysis of brain serotonin levels was performed using home made I-shaped probes, made of polyacrylonitrile / sodium methyl sulfonate copolymer dialysis fiber (i.d. 220 μ m, o.d. 0.31 μ m, AN 69, Hospal, Italy). Preceding surgery rats were anaesthetised using isoflurane (O₂/N₂O; 300/300ml/min). Lidocaine-HCl, 10 % (m/v) was used for local anaesthesia. Rats were placed in a stereotaxic frame (Kopf, USA), and probes were inserted into Ventral Hippocampus (V. Hippo, L +4.8 mm, IA: +3.7

mm, V: -8.0 mm) (Paxinos and Watson, 1982). After insertion, probes were secured with dental cement.

Microdialysis experiments

Rats were allowed to recover for at least 24 h. Probes were perfused with artificial cerebrospinal fluid containing 147 mM NaCl, 3.0 mM KCl, 1.2 mM CaCl_2 , and 1.2 mM MgCl_2 , at a flow-rate of 1.5 μl / min (Harvard apparatus, South Natick, Ma., USA). 15 minute microdialysis samples were collected in HPLC vials containing 7.5 μl 0.02 M acetic acid for serotonin analysis.

Serotonin analysis:

Twenty- μl microdialysate samples were injected via an autoinjector (CMA/200 refrigerated microsampler, CMA, Sweden) onto a 100 x 2.0 mm C18 Hypersil 3 μm column (Bester, Amstelveen, the Netherlands) and separated with a mobile phase consisting of 5 g/L di-ammoniumsulfate, 500 mg/L EDTA, 50 mg/L heptane sulphonic acid, 4 % methanol v/v, and 30 μl /L of triethylamine, pH 4.65 at a flow of 0.4 ml/min (Shimadzu LC-10 AD). 5-HT was detected amperometrically at a glassy carbon electrode at 500 mV vs Ag/AgCl (Antec Leyden, Leiden, The Netherlands). The detection limit was 0.5 fmol 5-HT per 20 μl sample (signal to noise ratio 3).

Data presentation and statistics

Four consecutive microdialysis samples with less than 20 % variation were taken as control and set at 100 %. Data are presented as percentages of control level (mean \pm S.E.M.) in time. Statistical analysis was performed using Sigmastat for Windows (SPSS, Jandel Corporation). Treatments were compared versus controls using two way analysis of variance (ANOVA) for repeated measurements, followed by Student Newman Keuls test. Drug effects were evaluated using one way ANOVA for repeated measures on ranks. Level of significance level was set at $p < 0.05$.

Results

Co-administration of citalopram with NFPS on 5-HT levels in ventral hippocampus

Administration of the glycine transporter inhibitor, NFPS (LU 2736N), at a dose of 10 $\mu\text{mol/kg}$ s.c. did not induce any significant effects on serotonin levels in rat ventral hippocampus ($X^2_{10} = 5.45$ $P = 0.857$). Co-administration of citalopram (10 $\mu\text{mol/kg}$ s.c.) together with the Glycine transporter inhibitor NFPS (10 $\mu\text{mol/kg}$ s.c.) significantly augmented the effect of citalopram on hippocampal serotonin levels (Treatment $F(1,9) = 5.35$, $P = 0.044$, Treatment vs. Time $F(1,104) = 2.12$, $P = 0.033$).

Claims:

1. The use of a compound, which is a serotonin reuptake inhibitor, and a compound, which is a GlyT-1 inhibitor for the preparation of a pharmaceutical composition for the treatment of depression, anxiety disorders and other affective disorders, such as generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder and social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse or any other disorder responsive to serotonin reuptake inhibitors.
2. The use of a GlyT-1 inhibitor for the preparation of a pharmaceutical composition to be used in combination with a serotonin reuptake inhibitor.
3. The use of a GlyT-1 inhibitor for the preparation of a pharmaceutical composition useful for augmenting and/or providing faster onset of the therapeutic effect of a serotonin reuptake inhibitor.
4. The use according to any of claims 2-3 wherein the serotonin reuptake inhibitor is used for the treatment of depression, anxiety disorders and other affective disorders, including generalized anxiety disorder, panic anxiety, obsessive compulsive disorder, acute stress disorder, post traumatic stress disorder or social anxiety disorder, eating disorders such as bulimia, anorexia and obesity, phobias, dysthymia, premenstrual syndrome, cognitive disorders, impulse control disorders, attention deficit hyperactivity disorder, drug abuse or any other disorder responsive to a SRI.
5. The use according to any one of claims 1-4 wherein the SRI is selected from a SSRI.
6. The use according to any one of claims 1-5 wherein the SRI is elected from citalopram, escitalopram, fluoxetine, sertraline, paroxetine, fluvoxamine, venlafaxine,

dapoxetine, duloxetine, vilazodone, nefazodone, imipramin, femoxetine and clomipramine.

7. The use according to any one of claims 1-6 wherein the GlyT-1 inhibitor is selected from

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl} glycine ethyl ester,

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine ethyl ester,

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl} glycine,

N-{3-[5-Cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[1-(3-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[1-(3-trifluoromethylphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[1-(3-trifluoromethylphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methyl (1-ethyl)glycine,

N-{3-[1-(4-methylphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylalanine,

N-{3-[1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methyl (1-ethyl)glycine,

N-{3-[4-chloro-1-(3-methyl-4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[4-chloro-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,

N-{3-[5-chloro-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylalanine,

N-{3-[6-chloro-1-(3-methyl-4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[6-chloro-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[6-chloro-1-(4-methylphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[6-chloro-1-(4-methoxyphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-fluoro-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-fluoro-1-(4-methoxyphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-trifluoromethyl-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-trifluoromethyl-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylalanine,
 N-{3-[5-cyano-1-(3-methyl-4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-cyano-1-(4-cyanophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylalanine,
 N-{3-[5-cyano-1-(4-methoxyphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-[5-cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine, N-{2-[5-cyano-1-(4-fluorophenyl)-1,3-dihydroisobenzofuran-1-yl]ethyl}-N-methylglycine,
 N-{3-[5-Chloro-1-(4-chloro-phenyl)-indan-1-yl]-propyl}-N-methylglycine,
 N-{3-[5-Chloro-1-(4-chloro-phenyl)-indan-1-yl]-propyl}-N-methylalanine,
 N-{3-[3-cyclo-1-(4-methylphenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl}-N-methylglycine,
 N-{3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-benzo[c]thiophen-1-yl)-propyl}-N-methylglycine,
 N-{3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-benzo[c]thiophen-1-yl)-propyl}-N-methylalanine,

N-[3-[1-(4-Fluoro-phenyl)-3,3-dimethyl-1,3-dihydro-isobenzofuran-1-yl]-propyl]-N-methylglycine,
 N-[3-[5-Bromo-1-(4-chlorophenyl)-1,3-dihydroisobenzofuran-1-yl]-1-propyl]-N-methylglycine,
 N-[2-[1-(4-Chloro-phenyl)-3,3-dimethyl-1,3-dihydro-isobenzofuran-1-yl]-ethyl]-N-methylglycine,
 N-[3-(3-methyl-1-phenyl-1*H*-inden-1-yl)-propyl]-N-methylglycine,
 N-[3-(5-Chloro-1-thiophen-2-yl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylglycine,
 N-[3-(5-Chloro-1-thiophen-2-yl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methyl (1-ethyl)-glycine,
 N-[3-(3-methyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylalanine,
 N-[3-(3-methyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methyl (1-ethyl)-glycine,
 N-[3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methylalanine,
 N-[3-(3,3-Dimethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methylalanine,
 N-[3-(3,3-Dimethyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methyl-(1-ethyl)glycine,
 N-[3-(3,3-Dimethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-ethyl]-N-methyl-(1-ethyl)glycine,
 N-[3-(3,3-Diethyl-1-phenyl-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylalanine,
 N-[3-(3,3-Diethyl-1-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylalanine,
 N-[3-(3,3-Diethyl-1-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl)-propyl]-N-methylglycine,
 N-[3-(1-phenyl-1,3-dihydro-benzo[c]thiophen-1-yl)-propyl]-N-methylalanine,
 N-[3-[1-(4-Chloro-phenyl)-3,3-dimethyl-indan-1-yl]-propyl]-N-methylglycine,
 N-[3-[1-(4-Chloro-phenyl)-3,3-diethyl-1,3-dihydro-isobenzofuran-1-yl]-propyl]-N-methyl-alanine,
 N-[2-(3-methyl-1-phenyl-indan-1-yl)-ethyl]-amino}-N-methyl alanine,

•

N-{3-[1-(4-Chloro-phenyl)-5-(3,4-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-ethyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(4-chloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(3-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2-methyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(2,5-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-Chloro-phenyl)-5-(3,4-dichloro-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine,
 N-{3-[1-(4-chloro-phenyl)-5-(2-trifluoromethyl-phenyl)-1,3-dihydro-isobenzofuran-1-yl]-propyl}-N-methyl-glycine ,
 (+/-)-{4-[2-(4-Methoxy-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-Chloro-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-Fluoro-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-*iso*-Propyl-phenylsulfanyl)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
 (+/-)-2-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-*trans*-2,5- dimethylpiperazin-1-yl}-propionic acid,
 {4-[5-Chloro-2-(4-methoxy-phenylsulfanyl)-phenyl]-2(R)-methyl-piperazin-1-yl}-acetic acid,
 {4-[2-(4-Methoxy-phenylsulfanyl)-phenyl]-2(R),5(S)-dimethyl-piperazin-1-yl}-acetic acid,

{4-[5-Chloro-2-(4-methoxy-phenylsulfanyl)-phenyl]-2,2-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[5-Chloro-2-(4-trifluoromethyl-phenylsulfanyl)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
 {4-[5-Chloro-2-(3-methoxy-phenylsulfanyl)-phenyl]-2(R)-methyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-Phenyl-phenyloxy)-phenyl]-2-methyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-Methyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(4-*iso*-Propyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-{4-[2-(2,4-Dimethyl-phenylsulfanyl)-phenyl]-*trans*-2,5-dimethyl-piperazin-1-yl}-acetic acid,
 (+/-)-2-{4-[2-(4-*tert*-Butyl-phenylsulfanyl)-phenyl]-3-methylpiperazin-1-yl}-propionic acid,
 {4-[2-(4-Isopropyl-phenylsulfanyl)-phenyl]-piperazin-1-yl}-acetic acid,
 (+/-)-2-{4-[2-(4-Methoxy-phenylsulfanyl)-phenyl]-3-methyl-piperazin-1-yl}-propionic acid,
 or a pharmaceutically acceptable acid addition salt thereof.

8. A pharmaceutical composition comprising a compound, which is a serotonin reuptake inhibitor, and a compound, which is a GlyT-1 inhibitor, and optionally pharmaceutically acceptable carriers or diluents.

9. The pharmaceutical composition of claim 8 wherein the SRI is selected from a SSRI.

10. The pharmaceutical composition of claim 8 or 9 wherein the GlyT-1 inhibitor show inhibition below 20000 nM, such as below 10000 nM, as IC₅₀ in the "[³H]-Glycine uptake" test described herein.

11. The pharmaceutical composition of any one of claims 8-10 wherein the GlyT-1 inhibitor is selected from any one of the inhibitors of claim 7.

12. The pharmaceutical composition according to any one of claims 8-11 characterized in that the serotonin uptake inhibitor is selected from citalopram, escitalopram, fluoxetine, sertraline, paroxetine, fluvoxamine, venlafaxine, dapoxetine, duloxetine, vilazodone, nefazodone, imipramin, femoxetine and clomipramine.
13. The pharmaceutical composition according to any one of claims 8-12, which is adapted for simultaneous administration of the active ingredients.
14. The pharmaceutical composition according to claim 13 wherein the active ingredients are contained in the same unit dosage form.
15. The pharmaceutical composition according to any one of claims 8-12 which is adapted for sequential administration of the active ingredients.
16. The pharmaceutical composition according to any one of claims 13 and 15 wherein the active ingredients are contained in discrete dosage forms.
17. A kit comprising a compound, which is a serotonin reuptake inhibitor, and a compound, which is a GlyT-1 inhibitor, and optionally pharmaceutically acceptable carriers or diluents.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/DK04/000547

International filing date: 18 August 2004 (18.08.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/496,738
Filing date: 21 August 2003 (21.08.2003)

Date of receipt at the International Bureau: 05 October 2004 (05.10.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.